



## FIRST AID EQUIPMENT

w. [www.first-aid-online.co.uk](http://www.first-aid-online.co.uk)

t. 01206 809 538

e. [support@protrainings.uk](mailto:support@protrainings.uk)



## TRAINING

w. [www.protrainings.uk](http://www.protrainings.uk)

t. 01206 805 359

e. [support@protrainings.uk](mailto:support@protrainings.uk)



ProTrainings Europe Ltd • Unit 22 Westside Centre • London Road  
Stanway • Colchester • Essex CO3 8PH



# Catastrophic Bleeding and Trauma First Aid Guidance

**FIRST AID** online  
[www.first-aid-online.co.uk](http://www.first-aid-online.co.uk)

## HOW TO USE WoundClot™ Haemostatic Dressing to stop a bleed

For full details, removal instructions and warnings please refer to the manufacturer's instructions. Please refer to the instructions below to ensure successful application of WoundClot™

Identify / Expose the wound

Identify the point of bleeding

Wipe away excess blood where possible.

Remove WoundClot™ from packet



If required, cut or tear WoundClot™ to the necessary size to suit the wound then place directly onto the wound. WoundClot™ can be formed to 'fit' any wound or site where rapid coagulation is required. Pack the wound with WoundClot™ if required.

Cover WoundClot™ with an appropriate dressing and apply direct pressure until bleeding is controlled /stops.

Wrap with a bandage to maintain pressure and seek further medical assistance.

WoundClot™ is available direct from ProTrainings [www.first-aid-online.co.uk](http://www.first-aid-online.co.uk)



## IMPORTANT Points to remember

### SERIOUS BLEEDING

Remember to wear gloves. To control bleeding apply pressure to the wound, apply a pressure dressing directly over the wound. If blood comes through dressing, remove and apply another dressing. Once bleeding controlled, activate EMS if needed or transport to medical help

### CATASTROPHIC BLEEDING - TOURNIQUETS

With limbs, you can apply a tourniquet tightly above the injury site but not over a joint till the bleeding stops. If required, apply another above the first. Note the time it was applied and tell EMS.

### CATASTROPHIC BLEEDING - HAEMOSTATIC DRESSING

Clear worst of blood to identify source. Apply Haemostatic dressing into the wound, apply pressure for required time. Apply trauma dressing and note time applied. Tuck wrapper under bandage for EMS.

**WoundClot can be used for all types of bleeds**

## CELOX GAUZE How to use it

### PACK

Pack any cavity with Celox Gauze.

### PRESS

Compress the wound to push the Celox Gauze on the area where the blood is coming from and stop blood flow for a short period while the Celox plug develops and strengthens.

### COMPRESS

Compress for 3 minutes or 1 minute with Celox Rapid

Call EMS and extract asap to medical services.



## TRAINING from ProTrainings

At ProTrainings we offer over 350 different subjects as video online, blended, virtual and nationwide classroom courses.

You can see our full range at [www.protrainings.uk/courses](http://www.protrainings.uk/courses)

For more information on any type of First Aid, Health & Safety, Medical, Care, HR, Social Care and much much more c ontact us on **01206 805359** or email [support@protrainings.uk](mailto:support@protrainings.uk) or you can chat to us online [www.protrainings.uk](http://www.protrainings.uk)

